

Antioxidative effects of alpha-lipoic acid in spinal cord injury

An experimental rat model



Ann. Ital. Chir., 2021 92, 1: 98-102
pii: S0003469X2103102X

Serdar Ercan*, Ayfer Aktaş**, Mustafa Serdar Kemalolu*

Dicle University, Medical Faculty, Diyarbakir, Turkey

*Department of Neurosurgery

**Department of Histology & Embryology

Antioxidative effects of alpha-lipoic acid in spinal cord injury. An experimental rat model

BACKGROUND: Traumatic and ischemic injuries of the spinal cord are effective in the development of neurological dysfunction of tissue damage caused by primary and secondary mechanisms. Free radical changes are effective in the development of early ischemia and progressive tissue ischemia is the main cause of secondary damage. Delaying ischemia is the basis of treatment. In this study, we aimed to demonstrate the presence of neuroprotective effects of alpha-lipoic acid in comparison with methylprednisolone.

METHODS: 50 Sprague Dawley rats were divided into 5 groups (n = 10) and spinal cord trauma was created by the method, described by Rivlin and Tator. Group 1: Laminectomy group, Group 2: Laminectomy + spinal cord injury (SCI), Group 3: Laminectomy + SCI + alpha-lipoic acid (ALA) (100 mg / kg), Group 4: Laminectomy + SCI + Methyl-prednisolone (30 mg / kg), Group 5: Laminectomy + SCI + ALA + Methyl-prednisolone.

RESULTS AND DISCUSSION: Rats with spinal cord injury were found to be paraplegic. There was no significant change in motor function between the groups. When the antioxidant values were compared in the groups, there was a statistically significant difference between Group 2 and Group 3. Oxygen radicals decreased significantly between ALA and Methylprednisolone. The most striking difference was between the monotherapy group and the combined treatment group.

CONCLUSION: Our results showed that alpha lipoic acid given after spinal cord trauma in rats decreases anti-oxidant formation.

KEY WORDS: Alpha-lipoic acid, Methyl-prednisolone, Oxygen radicals, Spinal cord injury

Introduction

Spinal cord injury, whether complete or incomplete, is a major social problem causes loss of major functions of the cord (motor, sensory, autonomic and reflex) ¹. It is hypothesized that there are two mechanisms for the damage caused by acute spinal cord injury. Primary injury resulting from mechanical tissue destruction results in necrotic cell death. Secondary damage, a cascade of

events triggered by injury, is the result of activation of endogenous cell death pathways ²⁻⁴. The majority of experimental and clinical studies in neuronal injury are aimed at preventing or reducing secondary injury ⁵. High-dose steroids are widely used in the treatment of spinal cord injury targeting many secondary degeneration mediators. But the results are not satisfactory because secondary degeneration mechanisms are not well understood ^{2,6,7}. Excitotoxicity, oxidative damage, ischemia, secondary injury are the contents of the cascade ^{5,8,9}.

The prominent factors in the occurrence of secondary injury is energy deficiency due to ischemia ^{9,10}. Theories of ischemia have been suggested as a decrease in tissue pH due to lactic acidosis, venous stasis and congestion due to fibrin and platelet accumulation, capillary endothelial damage, edema, petechial hemorrhages, and presence of vasoactive agents. Ischemia causes insufficient

Pervenuto in Redazione Giugno 2019. Accettato per la pubblicazione Agosto 2019.

Correspondence to: Serdar Ercan, Dicle University, Medical Faculty, Department of Neurosurgery, Yukari kilictasi mah, Sur, Diyarbakir, Turkey (e-mail: srdreercl@gmail.com)

glucose and oxygen intake to tissues and indirectly leads to a lack of energy and a decrease in ATP stores^{1,11}. Ischemia initiates the cascade of secondary pathogenetic mechanisms defined as excitotoxicity due to its dependence on endogenous excitatory amino acid neurotransmitters^{9,12,13}. In traumatic injury tissue repair, the inflammatory response begins, in which the activity of leukocytes is critical. Leukocytes remove cellular debris by phagocytosis, release enzymes that relax the extracellular matrix and secrete cytokines that direct proliferation of endothelial epithelial and connective tissue cells in the injured area^{14,15}. Cytokines accelerate the central nervous system (CNS) inflammatory response by inducing the expression of reactive oxygen and nitrogen derivatives of additional cytokines, chemokines, nitric oxide, inflammatory response¹⁶. When the first neural cell dies, it releases toxins that attack other living neurons. Some of these toxins are fugitive oxygen molecules called free radicals. Prevention of intensive free radical formation is vital for cell life¹⁷.

Alpha-lipoic acid is a substance synthesized naturally in our body and can be taken from food^{18,19}. There are two important functions in our body. The former acts as a coenzyme in the metabolic pathway, and the latter is an important antioxidant. In the experiments on rat model, alpha-lipoic acid enhances endoneurial blood flow, increases the physiological antioxidant level of glutathione and reduces free oxygen radicals in the diabetic nerve. These effects seen in the experimental state show that the function of peripheral nerves can be corrected with alpha-lipoic acid²⁰⁻²⁶.

Material and Methods

All experimental steps taken of this animal study is designed to eliminate pain and suffering with internationally accepted guidelines. Our experimental protocols were approved by University Animal Care and Use Committee.

50 Sprague Dawley rats with normal motor activity ranging between 200 and 250 g were included in the study. Rats were kept under standard laboratory conditions (12 hours day - 12 hours night lighting, 20-22 °C room temperature, 50-60% humidity). They were given enough water and food. They were randomly divided into 5 groups. Group 1: Laminectomy without any injury to spinal cord was performed. Group 2: Performed injury to the spinal cord but untreated. Group 3: Intramuscular Alpha-lipoic acid (ALA) was administered (100mg / kg) after laminectomy and nerve injury. Group 4: Intramuscular Methyl-prednisolone (30 mg / kg) was administered after laminectomy and nerve injury. Group 5: Combined treatment with ALA + Methyl-prednisolone after laminectomy and injury.

Injury: spinal cord injury was performed by the method, described by Rivlin and Tator²⁷. The rats were fasted

the day before and then anesthetized with Ketamine (60 mg/kg) + Xylazine (9 mg/kg) intramuscularly. The interscapular distance of the rats was referenced and a 3x2 cm area was shaved in the dorsal region and local antiseptics was applied with povidone iodine. Midline incision was performed at the T5-12 level in the prone position. The skin, subcutaneous tissues and paravertebral muscle fascia were passed through, and blunt dissection were implemented to the muscles for exploration of T7-10 laminae. Total laminectomy was performed at T7 level and extra care was taken not to damage the dura of the rats. At this level, the spinal cord was clamped with a Yasargil aneurysm clip (Aesculap FE 721K), which applied 63 g of force for standard trauma, for one minute to create spinal cord injury. The clip was then removed, and the anatomical layers were closed with 3/0 silk. The rats were placed in their cages. Bladders were emptied by manual compression twice daily and allowing free feeding. Treatments were administered according to group plans after the surgery.

Collecting the samples: 24 hours later intracardiac blood collected under Ketamine (60 mg / kg) + Xylazine (9 mg / kg) anesthesia After sacrifice the spinal cord was sectioned at the region of injury. Plasma samples obtained by rapid centrifugation from blood samples were transferred to plastic eppendorf cap tubes for biochemical analysis. Total antioxidant status (TAS), Total oxidant status (TOS), nitric oxide (NO), paraoxonase (PONX) activities were evaluated in samples by the same biochemist who was blind to groups.

Evaluation of functional recovery: Physical examinations were performed regularly to evaluate the functional recovery of rats. Motor functions of rats were evaluated immediately by Drummond and Moore criteria 24 hours after surgery (137). 0: paraplegic, no motor function in the lower extremities, 1: Motor function in the lower extremity is weak, only weak motion against gravity, 2: Moderate lower extremity motor function, good against gravity but cannot pull legs under the body, 3: Motor function Very good, he can pull his legs under the body and jump, but not completely normal, 4: Normal motor function.

Statistical analyze: Differences between groups were analyzed by Mann-Whitney U test.

Results

No mortality was seen in the groups after surgical intervention. There was no wound infection. Motor function of all rats was evaluated according to Drummond-Moore criteria (Table I). There was no significant difference between the groups ($p > 0.005$). TAS, TOS and NO values were compared between Group 2 and Group 3 and noticed that Sildenafil significantly increased TAS and NO values and decreased TOS value (TAS $p = 0.002$, TOS $p = 0.002$, NO $p = 0.003$). But there was no significant

TABLE I - Motor functions of the rats scored after surgery (Drummon – Moore score: 1. Motor function in the lower extremity is weak, only weak motion against gravity, 2. Moderate lower extremity motor function, good against gravity but cannot pull legs under the body, 3. Motor function Very good, he can pull his legs under the body and jump, 4. Normal motor function)

	Motor functions of the rats scored after surgery									
	Rat 1	Rat 2	Rat 3	Rat 4	Rat 5	Rat 6	Rat 7	Rat 8	Rat 9	Rat 10
Grup 1	4	4	4	4	3	4	4	3	4	4
Grup 2	1	1	1	1	2	1	1	2	1	1
Grup 3	2	2	1	0	0	1	1	2	0	0
Grup 4	2	2	1	0	0	0	0	0	1	1
Grup 5	2	2	1	0	2	2	1	1	0	0




Fig. 1: TAS, TOS, NO and PONX value of groups. (TAS: Total antioxidant status, TOS: Total oxidant status, NO: nitric oxide, PONX: paraoxonase)

difference in PONX values ($p = 0.221$). The similar results were repeated between the group 2 and 4, treated with methyl-prednisolone. There was no significant difference in PONX and NO values (PONX $p = 0.157$, NO $p = 0.941$), TAS and TOS values were similar to Sildenafil treated group (TAS $p = 0.002$, TOS $p = 0.034$). While the difference between group 2 and Sildenafil + Methyl-prednisolone combined treatment group (group 5) was similar in terms of TAS and TOS (TAS $p = 0.002$, TOS $p = 0.002$), NO values were significantly increased with sildenafil effect ($p = 0.003$). When group 3 and group 4 were compared, it was observed that sildenafil significantly increased TAS ($p = 0.01$), decreased TOS ($p = 0.001$), but did not make a significant change in NO and PONX (NO $p = 0.103$, PONX $p = 0.701$). The significant effect was seen in the combined treatment group. In the comparison of Group 3 and Group 5, TAS, TOS and NO values were significantly different (TAS $p = 0.031$, TOS $p = 0.002$, NO $p = 0.002$), PONX was not significant in this group ($p = 0.597$) (Fig. 1).

Discussion

After Rivlin and Tator described the myelotomy model with experimental spinal cord contusion, various treatment

approaches for spinal cord injury were presented²⁷. Many studies have been performed on the pathophysiology of spinal cord injury. Traumatic injuries of the spinal cord can cause primary and secondary tissue damage, which is also named direct and indirect damage. Experimental studies over the past twenty years have shown that these secondary mechanisms play a major role in the development of neurological dysfunction in spinal cord injury.

Acute treatments after spinal cord trauma are performed with neuroprotective drugs such as antioxidants, neurotransmitter receptor blockers, phosphokinase stimulators, phosphatase inhibitors. In the literature, many treatment methods for spinal cord injuries after experimental trauma have been investigated. In particular, Glutamate, Na channel blockers, erythropoietin, minocycline, erythropoietin, opiate antagonists, Ca channel blockers, gangliosides, amino acids, NOS, interleukin, lipopolysaccharides have been studied in many studies, but it is concluded that these treatments are healing enhancing treatments instead of direct treatment of the damage^{3,28}. Lipoic acid is an important coenzyme and antioxidant in the mechanism of pyruvate dehydrogenase and alpha ketoglutarate dehydrogenase. Lipoic acid has superiority over other antioxidants due to its water and oil solubility properties and it has been used as a treatment

in many diseases and new studies are still continuing. It has been used in the treatment of neurodegenerative diseases such as diabetic neuropathy, atherosclerosis and MS, joint disorders and infections like AIDS^{1,10,29,30}. In clinical studies, treatment of polyneuropathy and mononeuropathy due to diabetes, alpha lipoic acid has been shown to reduce neuropathic complaints of patients^{1,29,31}.

Clinical studies show that the spinal cord lesion is enlarged by secondary injury. The underlying molecular and cellular mechanisms are not fully understood. Current evidence suggests that free oxygen radical formation and peroxidation of membrane lipids play a role. The efficacy of some therapeutic approaches has been evaluated experimentally by determining the level of lipid peroxidation^{18,19,32-36}. Peroxidation has been evaluated by measuring the amount of malondialdehyde (MAD), one of the lipid peroxidation products, in the tissue^{16,18,32,33}. Although naturally occurring antioxidant compounds control harmful effects in normal cells, excessive free radical production in pathological conditions such as spinal cord injury exceeds antioxidant capacity. Free radicals lead to lipid peroxidation and, consequently, cell membrane dissolution. Antioxidants in the central nervous system are ascorbate, glutathione and alpha tocopherol. These antioxidants neutralize free radicals. Trauma reduces these substances^{33,37}.

Prevention of intensive free radical formation is an important first vital step for cell life. Methylprednisolone is one of the most important treatment protocols used to prevent the formation of free radicals. Other agents used are Glutamate, Na channel blockers, hypothermia, erythropoietin, minocycline, erythropoietin, opiate antagonists, Ca channel blockers, gangliosides, amino acids, NOS, interleukin, lipopolysaccharides. None of these substances used alone has been shown to be therapeutic, but it was found to be effective in combined use^{18,19,34-36,38-40}. The experiment was designed to compare antioxidative effect of alpha lipoic acid with Methylprednisolone which was commonly used in treatment of oxidative stress.

German researchers have found that lipoic acid has positive effects on long-term memory in older mice. Similar benefits have not yet been demonstrated in younger mice. Researchers have shown that lipoic acid compensates for long-term memory impairment rather than correcting general memory. Researchers at the University of Rochester medical center have reported a possible role in the treatment of chronic and acute neurological disorders such as Parkinson's and Huntington's disease¹³.

In conclusion, oxidative stress is the prime element in secondary injury of the spinal cord trauma. Avoid of intensive free radicals is crucial and antioxidant treatment should be used.

Alpha lipoic acid given after spinal cord trauma in rats increased the anti-oxidant formation and could be a treatment option on the patient who suffer from spinal

cord trauma or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript followed.

Riassunto

Le lesioni traumatiche e ischemiche del midollo spinale sono causa di disfunzione neurologica da danno tissutale per meccanismi primari e secondari. I cambiamenti dei radicali liberi sono dannosi per lo sviluppo dell'ischemia precoce e l'ischemia progressiva dei tessuti è la principale causa di danno secondario. Ritardare l'ischemia è la base del trattamento. In questo studio, abbiamo mirato a dimostrare la presenza di effetti neuroprotettivi dell'acido alfa-lipoico rispetto al metilprednisolone.

Sono stati impiegati sperimentalmente 50 ratti Sprague Dawley, divisi in 5 gruppi (ciascuno di 10 unità) e il trauma del midollo spinale è stato creato con il metodo, descritto da Rivlin e Tator. Gruppo 1: gruppo Laminectomia, Gruppo 2: Laminectomia + lesione del midollo spinale (SCI), Gruppo 3: Laminectomia + SCI + acido alfa-lipoico (ALA) (100 mg / kg), Gruppo 4: Laminectomia + SCI + Metil-prednisolone (30 mg / kg), Gruppo 5: Laminectomia + SCI + ALA + Metil-prednisolone.

RISULTATI E DISCUSSIONE: I ratti con lesione del midollo spinale sono risultati paraplegici. Non vi è stato alcun cambiamento significativo nella funzione motoria tra i gruppi. Quando i valori di antiossidanti sono stati confrontati nei gruppi, c'era una differenza statisticamente significativa tra il gruppo 2 e il gruppo 3. I radicali dell'ossigeno sono diminuiti significativamente tra ALA e metilprednisolone. La differenza più evidente era tra il gruppo di monoterapia e il gruppo di trattamento combinato.

In conclusione questi risultati hanno indicato che l'acido alfa lipoico somministrato dopo un trauma del midollo spinale nei ratti diminuisce la formazione di antiossidanti.

References

1. Dumont RJ, et al.: *Acute spinal cord injury, part II: contemporary pharmacotherapy*. Clin. Neuropharmacol, 2001; 24, 265-79.
2. Li M., et al.: *Functional role and therapeutic implications of neuronal caspase-1 and-3 in a mouse model of traumatic spinal cord injury*. Neuroscience, 2000; 99, 333-42.
3. Tator CH: *Strategies for recovery and regeneration after brain and spinal cord injury*. Inj Prev, 2002; 8:iv33-iv36.
4. Anderson TE: *Spinal cord contusion injury: experimental dissociation of hemorrhagic necrosis and subacute loss of axonal conduction*. J. Neurosurg, 1985; 62:115-19.
5. Hall ED, Wolf DL: *A pharmacological analysis of the pathophysiology of spinal cord injury*. J. Neurosurg, 1985; 62:115-19.

- biological mechanisms of posttraumatic spinal cord ischemia. J Neurosurg, 1986; 64:951-61.
6. Kirshblum SC, Groah SL, McKinley WO, Gittler MS, Stiens SA: *Etiology, classification, and acute medical management*. Arch Phys Med Rehabil, 2002; 83, S50-S57.
7. Anderson DK, Means ED, Waters TR, Green ES: *Microvascular perfusion and metabolism in injured spinal cord after methylprednisolone treatment*. J Neurosurg, 1982; 56, 106-13.
8. Faden AI, Simon R: *A potential role for excitotoxins in the pathophysiology of spinal cord injury*. Ann Neurol Off J Am Neurol Assoc Child Neurol Soc, 1988; 23: 623-26.
9. Park E, Velumian AA, Fehlings MG: *The role of excitotoxicity in secondary mechanisms of spinal cord injury: A review with an emphasis on the implications for white matter degeneration*. J Neurotrauma, 2004; 21:75-84.
10. Agrawa SK, Nashm R, Fehlings MG: *Role of L-and N-type calcium channels in the pathophysiology of traumatic spinal cord white matter injury*. Neuroscience, 2000; 99:179-88.
11. Tator CH, Fehlings MG: *Review of the secondary injury theory of acute spinal cord trauma with emphasis on vascular mechanisms*. J Neurosurg, 1991; 75:15-26.
12. Amar AP, Levy ML: *Pathogenesis and pharmacological strategies for mitigating secondary damage in acute spinal cord injury*. Neurosurgery, 1999; 44; 1027-39.
13. Oyar EÖ, Korkmaz A, Kardesş Ö, Ömeroğlu S: *Aortic cross-clamping-induced spinal cord oxidative stress in rabbits: The role of a novel antioxidant adrenomedullin*. J Surg Res, 2008; 147: 143-47.
14. Klusman I, Schwab ME: *Effects of pro-inflammatory cytokines in experimental spinal cord injury*. Brain Res, 7, 1997; 62, 173-84.
15. Zhang, Krebs C J, Gut, L: *Experimental analysis of progressive necrosis after spinal cord trauma in the rat: Etiological role of the inflammatory response*. Exp Neurol, 1997; 143:141-52.
16. Bethea JR, Dietrich DW: *Targeting the host inflammatory response in traumatic spinal cord injury*. Curr Opin Neurol, 2002; 15, 355-60.
17. Heffner JE, Repine-54.
18. Packe, L: *New horizons in antioxidant research: Action of the thioctic acid/dihydrolipoic acid couple in biological systems*. in Thioctacide. 2nd International Thioctic Acid Workshop, 1992; 35-44.
19. Packer L, Tritschle, H Wessel K: *Neuroprotection by the metabolic antioxidant α -lipoic acid*. Free Radic Biol Med, 1997; 22: 359-78.
20. Tankov, Cherninkova T, Koev D: *Treatment for diabetic mononeuropathy with α -lipoic acid*. Int J Clin Pract, 2005; 59, 645-50.
21. Fehlings MG: *Summary statement: the use of methylprednisolone in acute spinal cord injury*. Spine (Phila. Pa. 1976). 26, S55 (2001).
22. Genovese T. et al.: *Modulation of nitric oxide homeostasis in a mouse model of spinal cord injury*. J Neurosurg Spine, 2006; 4:145-53.
23. Cotter MA, Cameron NE, Greaves M: *The effects of treatment with [alpha]-lipoic acid or evening primrose oil on vascular hemostatic and lipid risk factors, blood flow, and peripheral nerve conduction in the streptozotocin-diabetic rat*. Metab Exp, 2001; 50:868-75 .
24. Şehirli AÖ, et al.: *Protective effects of alpha-lipoic acid against oxidative injury in tms-induced colitis*. Erciyes Med Journal/Erciyes Tip Derg, 2009; 31.
25. Toklu HZ, et al.: *The protective effect of alpha lipoic acid against traumatic brain injury in rats*. Free Radic Res, 43: 658-67.
26. Marracci G, Jones R, GM: *Alpha lipoic acid inhibits T cell migration into the spinal cord and suppresses and treats experimental autoimmune encephalomyelitis*. Elsevier, 2002.
27. Rivlin AS, Tator CH: *Effect of duration of acute spinal cord compression in a new acute cord injury model in the rat*. Surg Neurol, 1978; 10:38-43.
28. Schwab ME: *Repairing the injured spinal cord*. Science (80). 295, 2002; 1029-031.
29. Eaton MJ: *Cell and molecular approaches to the attenuation of pain after spinal cord injury*. J Neurotrauma, 2006; 23, 549-59.
30. Ellison D, et al.: *Neuropathology E-book: a reference text of CNS pathology*. Elsevier Health Sciences, 2012.
31. Ametov AS, et al.: *The sensory symptoms of diabetic polyneuropathy are improved with α -lipoic acid*. Diabetes Care, 2003; 26, 770-76.
32. Sekhon L, Fehlings HS: *Epidemiology, demographics, and pathophysiology of acute spinal cord injury*. Spine (Phila. Pa. 1976). 26, S2-S12 (2001).
33. Çakir E, Baykal S, Karahan CS, Kuzeyle K, Uydu H: *Acute phase effects of ATP-MgCl₂ on experimental spinal cord injury*. Neurosurg Rev, 2003; 26:67-70.
34. Delamarter RB, Sherman J, Carr JB: *Pathophysiology of spinal cord injury. Recovery after immediate and delayed decompression*. JBJS, 1995; 77:1042-049.
35. Díaz-Ruiz A, et al.: *Constitutive and inducible nitric oxide synthase activities after spinal cord contusion in rats*. Neurosci Lett, 2002; 319: 129-32.
36. Disse BG, Ziegler HW: *Pharmacodynamic mechanism and therapeutic activity of ambroxol in animal experiments*. Respiration, 1987; 51:15-22.
37. Klusmann S, Martin-Villalba A: *Molecular targets in spinal cord injury*. J Mol Med, 2005; 83:657-71.
38. Hall ED, Wolf DL, Braughler JM: *Effects of a single large dose of methylprednisolone sodium succinate on experimental posttraumatic spinal cord ischemia: Dose-response and time-action analysis*. J Neurosurg, 1984; 61, 124-30.
39. Ducke TB, Kind GW, Kempe LG: *Pathological findings in acute experimental spinal cord trauma*. J Neurosurg, 1971; 35, 700-08.
40. Duh MS, Shepar MJ, Wilberger JE, Bracken MB: *The effectiveness of surgery on the treatment of acute spinal cord injury and its relation to pharmacological treatment*. Neurosurgery, 1994; 35:240-49.